

Package ‘truncreg’

July 2, 2014

Version 0.2-1

Date 2013-12-24

Title Truncated Gaussian Regression Models

Depends R (>= 1.8.0), maxLik

Suggests survival

Description Estimation of models for truncated Gaussian variables by maximum likelihood.

License GPL (>= 2)

URL <http://R-Forge.R-project.org/projects/truncreg/>

Author Yves Croissant [aut, cre], Achim Zeileis [aut]

Maintainer Yves Croissant <yves.croissant@univ-reunion.fr>

Repository CRAN

Repository/R-Forge/Project truncreg

Repository/R-Forge/Revision 5

Repository/R-Forge/DateTimeStamp 2013-12-24 09:59:06

Date/Publication 2013-12-27 13:28:25

NeedsCompilation no

R topics documented:

truncreg	2
Index	5

truncreg

*Truncated Gaussian Response Models***Description**

Estimation of models for truncated Gaussian variables by maximum likelihood.

Usage

```
truncreg(formula, data, subset, weights, na.action,
  point = 0, direction = "left",
  model = TRUE, y = FALSE, x = FALSE, ...)
```

Arguments

formula	a symbolic description of the model to be estimated,
data	the data,
subset	an optional vector specifying a subset of observations,
weights	an optional vector of weights,
na.action	a function which indicates what should happen when the data contains 'NA's,
point	the value of truncation (the default is 0),
direction	the direction of the truncation, either "left" (the default) or "right",
model, y, x	logicals. If TRUE the corresponding components of the fit (model frame, response, model matrix) are returned.
...	further arguments.

Details

The model is estimated with the [maxLik](#) package and the Newton-Raphson method, using analytic gradient and Hessian.

A set of standard extractor functions for fitted model objects is available for objects of class "truncreg", including methods to the generic functions [print](#), [summary](#), [coef](#), [vcov](#), [logLik](#), [residuals](#), [predict](#), [fitted](#), [model.frame](#), and [model.matrix](#).

Value

An object of class "truncreg", a list with elements:

coefficients	the named vector of coefficients,
vcov	the variance matrix of the coefficients,
fitted.values	the fitted values,
logLik	the value of the log-likelihood,
gradient	the gradient of the log-likelihood at convergence,

nobs	the number of observations,
call	the matched call,
terms	the model terms,
model	the model frame used (if model = TRUE),
y	the response vector (if y = TRUE),
x	the model matrix (if x = TRUE),
point	the truncation point used,
direction	the truncation direction used,
est.stat	some information about the estimation (time used, optimization method),

References

Cragg JG (1971). Some Statistical Models for Limited Dependent Variables with Application to the Demand for Durable Goods. *Econometrica*, **39**, 829–844.

Hausman JA, Wise DA (1976). The Evaluation of Results from Truncated Samples: The New-Jersey Negative Income Tax Experiment. *Annals of Economic and Social Measurement*, **5**, 421–445.

Hausman JA, Wise DA (1976). Social Experimentation, Truncated Distributions and Efficient Estimation. *Econometrica*, **45**, 421–425.

Tobin J (1958). Estimation of Relationships for Limited Dependent Variables. *Econometrica*, **26**, 24–36.

See Also

[maxLik](#), [mhurdle](#)

Examples

```
#####
## Artificial example ##
#####

## simulate a data.frame
set.seed(1071)
n <- 10000
sigma <- 4
alpha <- 2
beta <- 1
x <- rnorm(n, mean = 0, sd = 2)
eps <- rnorm(n, sd = sigma)
y <- alpha + beta * x + eps
d <- data.frame(y = y, x = x)

## truncated response
d$yt <- ifelse(d$y > 1, d$y, NA)

## binary threshold response
d$yb <- factor(d$y > 0)
```

```

## censored response
d$yc <- pmax(1, d$y)

## compare estimates for full/truncated/censored/threshold response
fm_full <- lm(y ~ x, data = d)
fm_trunc <- truncreg(yt ~ x, data = d, point = 1, direction = "left")
fm_thresh <- glm(yb ~ x, data = d, family = binomial(link = "probit"))
library("survival")
fm_cens <- survreg(Surv(yc, yc > 1, type = "left") ~ x, data = d, dist = "gaussian")

## compare scaled regression coefficients
cbind(
  "True"      = c(alpha, beta) / sigma,
  "Full"      = coef(fm_full) / summary(fm_full)$sigma,
  "Truncated" = coef(fm_trunc)[1:2] / coef(fm_trunc)[3],
  "Censored"  = coef(fm_cens) / fm_cens$scale,
  "Threshold" = coef(fm_thresh)
)

#####
## Tobin's durable goods data ##
#####

## Tobit model (Tobin 1958)
data("tobin", package = "survival")
tobit <- survreg(Surv(durable, durable > 0, type = "left") ~ age + quant,
  data = tobin, dist = "gaussian")

## Two-part model (Cragg 1971)
## (see "mhurdle" package for a combined solution)
cragg_probit <- glm(factor(durable > 0) ~ age + quant,
  data = tobin, family = binomial(link = "logit"))
cragg_trunc <- truncreg(durable ~ age + quant, data = tobin, subset = durable > 0)

## Scaled coefficients
cbind(
  "Tobit"      = coef(tobit) / tobit$scale,
  "Binary"     = coef(cragg_probit),
  "Truncated"  = coef(cragg_trunc)[1:3] / coef(cragg_trunc)[4])

## likelihood ratio test and BIC
ll <- c("Tobit" = tobit$loglik[1],
  "Two-Part" = as.vector(logLik(cragg_probit) + logLik(cragg_trunc)))
df <- c(4, 3 + 4)
pchisq(2 * diff(ll), diff(df), lower.tail = FALSE)
-2 * ll + log(nrow(tobin)) * df

```

Index

*Topic **regression**

truncreg, 2

coef, 2

coef.truncreg (truncreg), 2

fitted, 2

fitted.truncreg (truncreg), 2

logLik, 2

logLik.truncreg (truncreg), 2

maxLik, 2, 3

mhurdle, 3

model.frame, 2

model.frame.truncreg (truncreg), 2

model.matrix, 2

model.matrix.truncreg (truncreg), 2

predict, 2

predict.truncreg (truncreg), 2

print, 2

print.summary.truncreg (truncreg), 2

print.truncreg (truncreg), 2

residuals, 2

residuals.truncreg (truncreg), 2

summary, 2

summary.truncreg (truncreg), 2

truncreg, 2

vcov, 2

vcov.truncreg (truncreg), 2