

Package ‘Peptides’

November 16, 2020

Version 2.4.3

Title Calculate Indices and Theoretical Physicochemical Properties of Protein Sequences

URL <https://github.com/dosorio/Peptides/>

Suggests testthat (>= 2.1.0)

Description Includes functions to calculate several physicochemical properties and indices for amino-acid sequences as well as to read and plot 'XVG' output files from the 'GROMACS' molecular dynamics package.

License GPL-2

LinkingTo Rcpp

Imports Rcpp

RoxygenNote 7.1.1

Encoding UTF-8

NeedsCompilation yes

Author Daniel Osorio [aut, cre] (<<https://orcid.org/0000-0003-4424-8422>>),
Paola Rondon-Villarreal [aut, ths] (<<https://orcid.org/0000-0001-8209-3885>>),
Rodrigo Torres [aut, ths] (<<https://orcid.org/0000-0003-1113-3020>>),
J. Sebastian Paez [ctb] (<<https://orcid.org/0000-0002-0065-1474>>),
Luis Pedro Coelho [ctb] (<<https://orcid.org/0000-0002-9280-7885>>),
Richèl J.C. Bilderbeek [ctb] (<<https://orcid.org/0000-0003-1107-7049>>),
Florian C. Sigloch [ctb] (<<https://orcid.org/0000-0001-8130-6885>>)

Maintainer Daniel Osorio <dcosorioh@tamu.edu>

Repository CRAN

Date/Publication 2020-11-16 17:10:02 UTC

R topics documented:

aaComp	2
AAdata	4
aaDescriptors	9

aaList	10
aaSMILES	11
aIndex	12
autoCorrelation	13
autoCovariance	14
blosumIndices	15
boman	16
charge	17
crossCovariance	18
crucianiProperties	20
fasgaiVectors	21
hmoment	22
hydrophobicity	23
instaIndex	27
kideraFactors	28
lengthpep	29
massShift	30
membpos	31
mswhimScores	32
mw	33
mz	34
pepdata	35
pI	36
plotXVG	37
protFP	38
readXVG	39
stScales	40
tScales	41
vhseScales	41
zScales	42
Index	44

aaComp

Compute the amino acid composition of a protein sequence

Description

This function calculates the amount of amino acids of a particular class and classified as: Tiny, Small, Aliphatic, Aromatic, Non-polar, Polar, Charged, Basic and Acidic based on their size and R-groups using same function implemented in EMBOSS 'pepstat'. The output is a matrix with the number and percentage of amino acids of a particular class

Usage

```
aaComp(seq)
```

Arguments

seq An amino-acid sequence

Details

Amino acids are zwitterionic molecules with an amine and a carboxyl group present in their structure. Some amino acids possess side chains with specific properties that allow grouping them in different ways. The aaComp function classifies amino acids based on their size, side chains, hydrophobicity, charge and their response to pH 7.

Value

The output is a matrix with the number and percentage of amino acids of a particular class

- Tiny (A + C + G + S + T)
- Small (A + B + C + D + G + N + P + S + T + V)
- Aliphatic (A + I + L + V)
- Aromatic (F + H + W + Y)
- Non-polar (A + C + F + G + I + L + M + P + V + W + Y)
- Polar (D + E + H + K + N + Q + R + S + T + Z)
- Charged (B + D + E + H + K + R + Z)
- Basic (H + K + R)
- Acidic (B + D + E + Z)

Note

This function was originally written by Alan Bleasby (ajb@ebi.ac.uk) for the EMBOSS package. Further information: <http://emboss.sourceforge.net/apps/cvs/emboss/apps/pepstats.html>

References

Rice, Peter, Ian Longden, and Alan Bleasby. "EMBOSS: the European molecular biology open software suite." Trends in genetics 16.6 (2000): 276-277.

Examples

```
# COMPARED TO PEPSTATS
# http://emboss.bioinformatics.nl/cgi-bin/emboss/pepstats
# Property      Residues          Number  Mole%
# Tiny          (A+C+G+S+T)      4       19.048
# Small         (A+B+C+D+G+N+P+S+T+V) 4       19.048
# Aliphatic     (A+I+L+V)        5       23.810
# Aromatic      (F+H+W+Y)        5       23.810
# Non-polar     (A+C+F+G+I+L+M+P+V+W+Y) 11      52.381
# Polar         (D+E+H+K+N+Q+R+S+T+Z) 9       42.857
# Charged       (B+D+E+H+K+R+Z)  8       38.095
# Basic         (H+K+R)          8       38.095
# Acidic        (B+D+E+Z)        0       00.000
```

```
## AA composition of PDB: 1D9J Cecropin Peptide
aaComp(seq= "KWKLFKKIGIGKFLHSAKKFX")
```

```
## Output
#           Number  Mole %
# Tiny           4 19.048
# Small          4 19.048
# Aliphatic      5 23.810
# Aromatic       5 23.810
# NonPolar      11 52.381
# Polar          9 42.857
# Charged        8 38.095
# Basic          8 38.095
# Acidic         0  0.000
```

AAdata	<i>Properties, scales and indices for the 20 naturally occurring amino acids from various sources</i>
--------	-------------------------------------------------------------------------------------------------------

Description

A list with a collection of properties, scales and indices for the 20 naturally occurring amino acids from various sources.

Usage

```
data(AAdata)
```

Format

A list as follows:

- **Hydrophobicity** The hydrophobicity is an important stabilization force in protein folding; this force changes depending on the solvent in which the protein is found.
 - **Aboderin**: Aboderin, A. A. (1971). An empirical hydrophobicity scale for alpha-amino-acids and some of its applications. *International Journal of Biochemistry*, 2(11), 537-544.
 - **AbrahamLeo**: Abraham D.J., Leo A.J. Hydrophobicity (delta G1/2 cal). *Proteins: Structure, Function and Genetics* 2:130-152(1987).
 - **Argos**: Argos, P., Rao, J. K., & Hargrave, P. A. (1982). Structural Prediction of Membrane-Bound Proteins. *European Journal of Biochemistry*, 128(2-3), 565-575.
 - **BlackMould**: Black S.D., Mould D.R. Hydrophobicity of physiological L-alpha amino acids. *Anal. Biochem.* 193:72-82(1991).
 - **BullBreese**: Bull H.B., Breese K. Hydrophobicity (free energy of transfer to surface in kcal/mole). *Arch. Biochem. Biophys.* 161:665-670(1974).
 - **Casari**: Casari, G., & Sippl, M. J. (1992). Structure-derived hydrophobic potential: hydrophobic potential derived from X-ray structures of globular proteins is able to identify native folds. *Journal of molecular biology*, 224(3), 725-732.

- Chothia: Chothia, C. (1976). The nature of the accessible and buried surfaces in proteins. *Journal of molecular biology*, 105(1), 1-12.
- Cid: Cid, H., Bunster, M., Canales, M., & Gazitua, F. (1992). Hydrophobicity and structural classes in proteins. *Protein engineering*, 5(5), 373-375.
- Cowan3.4: Cowan R., Whittaker R.G. Hydrophobicity indices at pH 3.4 determined by HPLC. *Peptide Research* 3:75-80(1990).
- Cowan7.5: Cowan R., Whittaker R.G. Hydrophobicity indices at pH 7.5 determined by HPLC. *Peptide Research* 3:75-80(1990).
- Eisenberg: Eisenberg D., Schwarz E., Komarony M., Wall R. Normalized consensus hydrophobicity scale. *J. Mol. Biol.* 179:125-142(1984).
- Engelman: Engelman, D. M., Steitz, T. A., & Goldman, A. (1986). Identifying nonpolar transbilayer helices in amino acid sequences of membrane proteins. *Annual review of biophysics and biophysical chemistry*, 15(1), 321-353.
- Fasman: Fasman, G. D. (Ed.). (1989). *Prediction of protein structure and the principles of protein conformation*. Springer.
- Fauchere: Fauchere J.-L., Pliska V.E. Hydrophobicity scale (pi-r). *Eur. J. Med. Chem.* 18:369-375(1983).
- Goldsack: Goldsack, D. E., & Chalifoux, R. C. (1973). Contribution of the free energy of mixing of hydrophobic side chains to the stability of the tertiary structure of proteins. *Journal of theoretical biology*, 39(3), 645-651.
- Guy: Guy H.R. Hydrophobicity scale based on free energy of transfer (kcal/mole). *Biophys J.* 47:61-70(1985).
- HoppWoods: Hopp T.P., Woods K.R. Hydrophilicity. *Proc. Natl. Acad. Sci. U.S.A.* 78:3824-3828(1981).
- Janin: Janin J. Free energy of transfer from inside to outside of a globular protein. *Nature* 277:491-492(1979).
- Jones: Jones, D. D. (1975). Amino acid properties and side-chain orientation in proteins: a cross correlation approach. *Journal of theoretical biology*, 50(1), 167-183.
- Juretic: Juretic, D., Lucic, B., Zucic, D., & Trinajstic, N. (1998). Protein transmembrane structure: recognition and prediction by using hydrophobicity scales through preference functions. *Theoretical and computational chemistry*, 5, 405-445.
- Kidera: Kidera, A., Konishi, Y., Oka, M., Ooi, T., & Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, 4(1), 23-55.
- Kuhn: Kuhn, L. A., Swanson, C. A., Pique, M. E., Tainer, J. A., & Getzoff, E. D. (1995). Atomic and residue hydrophilicity in the context of folded protein structures. *Proteins: Structure, Function, and Bioinformatics*, 23(4), 536-547.
- KyteDoolittle: Kyte J., Doolittle R.F. Hydrophobicity. *J. Mol. Biol.* 157:105-132(1982).
- Levitt: Levitt, M. (1976). A simplified representation of protein conformations for rapid simulation of protein folding. *Journal of molecular biology*, 104(1), 59-107.
- Manavalan: Manavalan P., Ponnuswamy Average surrounding hydrophobicity. *P.K. Nature* 275:673-674(1978).
- Miyazawa: Miyazawa S., Jernigen R.L. Hydrophobicity scale (contact energy derived from 3D data). *Macromolecules* 18:534-552(1985).
- Parker: Parker J.M.R., Guo D., Hodges R.S. Hydrophilicity scale derived from HPLC peptide retention times. *Biochemistry* 25:5425-5431(1986).

- Ponnuswamy: Ponnuswamy, P. K. (1993). Hydrophobic characteristics of folded proteins. *Progress in biophysics and molecular biology*, 59(1), 57-103.
- Prabhakaran: Prabhakaran, M. (1990). The distribution of physical, chemical and conformational properties in signal and nascent peptides. *Biochem. J*, 269, 691-696.
- Rao: Rao M.J.K., Argos P. Membrane buried helix parameter. *Biochim. Biophys. Acta* 869:197-214(1986).
- Rose: Rose G.D., Geselowitz A.R., Lesser G.J., Lee R.H., Zehfus M.H. Mean fractional area loss (f) [average area buried/standard state area]. *Science* 229:834-838(1985)
- Roseman: Roseman M.A. Hydrophobicity scale (pi-r). *J. Mol. Biol.* 200:513-522(1988).
- Sweet: Sweet R.M., Eisenberg D. Optimized matching hydrophobicity (OMH). *J. Mol. Biol.* 171:479-488(1983).
- Tanford: Tanford C. Hydrophobicity scale (Contribution of hydrophobic interactions to the stability of the globular conformation of proteins). *J. Am. Chem. Soc.* 84:4240-4274(1962).
- Welling: Welling G.W., Weijer W.J., Van der Zee R., Welling-Wester S. Antigenicity value X 10. *FEBS Lett.* 188:215-218(1985).
- Wilson: Wilson K.J., Honegger A., Stotzel R.P., Hughes G.J. Hydrophobic constants derived from HPLC peptide retention times. *Biochem. J.* 199:31-41(1981).
- Wolfenden: Wolfenden R.V., Andersson L., Cullis P.M., Southgate C.C.F. Hydration potential (kcal/mole) at 25C. *Biochemistry* 20:849-855(1981).
- Zimmerman: Zimmerman, J. M., Eliezer, N., & Simha, R. (1968). The characterization of amino acid sequences in proteins by statistical methods. *Journal of theoretical biology*, 21(2), 170-201.
- interfaceScale_pH8 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- interfaceScale_pH2 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- octanolScale_pH8 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- octanolScale_pH2 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- oiScale_pH8 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- oiScale_pH2 White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25
- crucialProperties: The three Cruciani et. al (2004) properties, are the scaled principal component scores that summarize a broad set of descriptors calculated based on the interaction of each amino acid residue with several chemical groups (or "probes"), such as charged ions, methyl, hydroxyl groups, and so forth.
 - PP1: Polarity,
 - PP2: Hydrophobicity,
 - PP3: H-bonding
- kideraFactors: The Kidera Factors were originally derived by applying multivariate analysis to 188 physical properties of the 20 amino acids and using dimension reduction techniques. A 10-dimensional vector of orthogonal factors was then obtained for each amino acid. The first

four factors are essentially pure physical properties; the remaining six factors are superpositions of several physical properties, and are labelled for convenience by the name of the most heavily weighted component

- helix.bend.pref: Helix/bend preference
 - side.chain.size: Side-chain size
 - extended.str.pref: Extended structure preference
 - hydrophobicity: Hydrophobicity
 - double.bend.pref: Double-bend preference
 - partial.spec.vol: Partial specific volume
 - flat.ext.pref: Flat extended preference
 - occurrence.alpha.reg: Occurrence in alpha region
 - pK.C: pK-C
 - surrounding.hydrop: Surrounding hydrophobicity
- pK
 - Bjellqvist: Bjellqvist, B., Hughes, G.J., Pasquali, Ch., Paquet, N., Ravier, F., Sanchez, J.Ch., Frutiger, S., Hochstrasser D. (1993) The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis*, 14:1023-1031.
 - Dawson: Dawson, R. M. C.; Elliot, D. C.; Elliot, W. H.; Jones, K. M. *Data for biochemical research*. Oxford University Press, 1989; p. 592.
 - EMBOSS: EMBOSS data are from <http://emboss.sourceforge.net/apps/release/5.0/emboss/apps/iep.html>.
 - Lehninger: Nelson, D. L.; Cox, M. M. *Lehninger Principles of Biochemistry*, Fourth Edition; W. H. Freeman, 2004; p. 1100.
 - Murray: Murray, R.K., Granner, D.K., Rodwell, V.W. (2006) *Harper's illustrated Biochemistry*. 27th edition. Published by The McGraw-Hill Companies.
 - Rodwell: Rodwell, J. Heterogeneity of component bands in isoelectric focusing patterns. *Analytical Biochemistry*, 1982, 119 (2), 440-449.
 - Sillero: Sillero, A., Maldonado, A. (2006) Isoelectric point determination of proteins and other macromolecules: oscillating method. *Comput Biol Med.*, 36:157-166.
 - Solomon: Solomon, T.W.G. (1998) *Fundamentals of Organic Chemistry*, 5th edition. Published by Wiley.
 - Stryer: Stryer L. (1999) *Biochemia*. czwarta edycja. Wydawnictwo Naukowe PWN.
 - zScales The five Sandberg et al. (1998) Z-scales describe each amino acid with numerical values, descriptors, which represent the physicochemical properties of the amino acids including NMR data and thin-layer chromatography (TLC) data.
 - Z1: Lipophilicity
 - Z2: Steric properties (Steric bulk/Polarizability)
 - Z3: Electronic properties (Polarity / Charge)
 - Z4: Related to electronegativity, heat of formation, electrophilicity and hardness.
 - Z5: Related to electronegativity, heat of formation, electrophilicity and hardness.
 - FASGAI Factor Analysis Scale of Generalized Amino Acid Information (FASGAI) proposed by Liang and Li (2007), is a set of amino acid descriptors, that reflects hydrophobicity, alpha and turn propensities, bulky properties, compositional characteristics, local flexibility, and electronic properties, was derived from multi-dimensional properties of 20 naturally occurring amino acids.

- F1: Hydrophobicity index
- F2: Alpha and turn propensities
- F3: Bulky properties
- F4: Compositional characteristic index
- F5: Local flexibility
- F6: Electronic properties
- VHSE The principal components score Vectors of Hydrophobic, Steric, and Electronic properties, is derived from principal components analysis (PCA) on independent families of 18 hydrophobic properties, 17 steric properties, and 15 electronic properties, respectively, which are included in total 50 physicochemical variables of 20 coded amino acids.
 - VHSE1 and VHSE2: Hydrophobic properties
 - VHSE3 and VHSE4: Steric properties
 - VHSE5 to VHSE8: Electronic properties

Source

- Hydrophobicity
 - ExPASy-ProtScale (<http://web.expasy.org/protscale/>)
 - AAIndex Database (<http://www.genome.jp/aaindex/>)
- pK
 - Kiraga, J. (2008) Analysis and computer simulations of variability of isoelectric point of proteins in the proteomes. PhD thesis, University of Wroclaw, Poland.

References

- Hydrophobicity
 - Nakai, K., Kidera, A., and Kanehisa, M.; Cluster analysis of amino acid indices for prediction of protein structure and function. *Protein Eng.* 2, 93-100 (1988).
 - Tomii, K. and Kanehisa, M.; Analysis of amino acid indices and mutation matrices for sequence comparison and structure prediction of proteins. *Protein Eng.* 9, 27-36 (1996).
 - Kawashima, S., Ogata, H., and Kanehisa, M.; AAindex: amino acid index database. *Nucleic Acids Res.* 27, 368-369 (1999).
 - Kawashima, S. and Kanehisa, M.; AAindex: amino acid index database. *Nucleic Acids Res.* 28, 374 (2000).
 - Kawashima, S., Pokarowski, P., Pokarowska, M., Kolinski, A., Katayama, T., and Kanehisa, M.; AAindex: amino acid index database, progress report 2008. *Nucleic Acids Res.* 36, D202-D205 (2008).
- crucialProperties:
 - Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. *J. Chemom.* 18, 146-155.
- kideraFactors:
 - Kidera, A., Konishi, Y., Oka, M., Ooi, T., & Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, 4(1), 23-55.

- pK:
 - Aronson, J. N. The Henderson-Hasselbalch equation revisited. *Biochemical Education*, 1983, 11 (2), 68.
 - Moore, D. S.. Amino acid and peptide net charges: A simple calculational procedure. *Biochemical Education*, 1986, 13 (1), 10-12.
 - Goloborodko, A.A.; Levitsky, L.I.; Ivanov, M.V.; and Gorshkov, M.V. (2013) "Pyteomics - a Python Framework for Exploratory Data Analysis and Rapid Software Prototyping in Proteomics", *Journal of The American Society for Mass Spectrometry*, 24(2), 301-304.
 - Kiraga, J. (2008) Analysis and computer simulations of variability of isoelectric point of proteins in the proteomes. PhD thesis, University of Wroclaw, Poland.
- zScales
 - Sandberg M, Eriksson L, Jonsson J, Sjostrom M, Wold S: New chemical descriptors relevant for the design of biologically active peptides. A multivariate characterization of 87 amino acids. *J Med Chem* 1998, 41:2481-2491.
- FASGAI
 - Liang, G., & Li, Z. (2007). Factor analysis scale of generalized amino acid information as the source of a new set of descriptors for elucidating the structure and activity relationships of cationic antimicrobial peptides. *Molecular Informatics*, 26(6), 754-763.
- VHSE
 - Mei, H. U., Liao, Z. H., Zhou, Y., & Li, S. Z. (2005). A new set of amino acid descriptors and its application in peptide QSARs. *Peptide Science*, 80(6), 775-786.

aaDescriptors

Compute 66 descriptors for each amino acid of a protein sequence.

Description

The function return 66 amino acid descriptors for the 20 natural amino acids. Available descriptors are:

- crucialProperties: Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. *J. Chemom.* 18, 146-155.,
- kideraFactors: Kidera, A., Konishi, Y., Oka, M., Ooi, T., & Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, 4(1), 23-55.,
- zScales: Sandberg M, Eriksson L, Jonsson J, Sjostrom M, Wold S: New chemical descriptors relevant for the design of biologically active peptides. A multivariate characterization of 87 amino acids. *J Med Chem* 1998, 41:2481-2491.,
- FASGAI: Liang, G., & Li, Z. (2007). Factor analysis scale of generalized amino acid information as the source of a new set of descriptors for elucidating the structure and activity relationships of cationic antimicrobial peptides. *Molecular Informatics*, 26(6), 754-763.,
- tScales: Tian F, Zhou P, Li Z: T-scale as a novel vector of topological descriptors for amino acids and its application in QSARs of peptides. *J Mol Struct.* 2007, 830: 106-115. 10.1016/j.molstruc.2006.07.004.,

- VHSE: VHSE-scales (principal components score Vectors of Hydrophobic, Steric, and Electronic properties), is derived from principal components analysis (PCA) on independent families of 18 hydrophobic properties, 17 steric properties, and 15 electronic properties, respectively, which are included in total 50 physicochemical variables of 20 coded amino acids.,
- protFP: van Westen, G. J., Swier, R. F., Wegner, J. K., IJzerman, A. P., van Vlijmen, H. W., & Bender, A. (2013). Benchmarking of protein descriptor sets in proteochemometric modeling (part 1): comparative study of 13 amino acid descriptor sets. *Journal of cheminformatics*, 5(1), 41.,
- stScales: Yang, L., Shu, M., Ma, K., Mei, H., Jiang, Y., & Li, Z. (2010). ST-scale as a novel amino acid descriptor and its application in QSAM of peptides and analogues. *Amino acids*, 38(3), 805-816.,
- BLOSUM: Georgiev, A. G. (2009). Interpretable numerical descriptors of amino acid space. *Journal of Computational Biology*, 16(5), 703-723.,
- MSWHIM: Zaliani, A., & Gancia, E. (1999). MS-WHIM scores for amino acids: a new 3D-description for peptide QSAR and QSPR studies. *Journal of chemical information and computer sciences*, 39(3), 525-533.

Usage

```
aaDescriptors(seq)
```

Arguments

seq An amino-acids sequence. If multiple sequences are given all of them must have the same length (gap symbols are allowed.)

Value

a matrix with 66 amino acid descriptors for each aminoacid in a protein sequence.

Examples

```
aaDescriptors(seq = "KLKLLLLLKLK")
```

```
aaList
```

Return a vector with the 20 standard amino acids in upper case

Description

This function returns a vector with the 20 standard amino acids in upper case.

Usage

```
aaList()
```

Value

A character vector with the 20 standard amino acids in upper case.

Author(s)

Richel Bilderbeek <richel@richelbilderbeek.nl>

References

Lu, Y., & Freeland, S. (2006). On the evolution of the standard amino-acid alphabet. *Genome biology*, 7(1), 102.

aaSMILES

Create Smiles String from aminoacid sequences

Description

This function converts peptides with aminoacid one-letter abbreviations into smiles strings to represent the structure.

Usage

```
aaSMILES(seq)
```

Arguments

seq character vector with one-letter aminoacid codes

Details

The output can be stored in a .smi file and converted using openbabel to drawings of the peptides.

Value

character vector with smiles strings

Examples

```
aaSMILES("AA")
# [1] "N[C@]([H])(C)C(=O)N[C@]([H])(C)C(=O)O"
aaSMILES(c("AA", "GG"))
# [1] "N[C@]([H])(C)C(=O)N[C@]([H])(C)C(=O)O" "NCC(=O)NCC(=O)O"
```

`aIndex`*Compute the aliphatic index of a protein sequence*

Description

This function calculates the Ikai (1980) aliphatic index of a protein. The `aIndex` is defined as the relative volume occupied by aliphatic side chains (Alanine, Valine, Isoleucine, and Leucine). It may be regarded as a positive factor for the increase of thermostability of globular proteins.

Usage

```
aIndex(seq)
```

Arguments

```
seq          An amino-acids sequence
```

Details

Aliphatic amino acids (A, I, L and V) are responsible for the thermal stability of proteins. The aliphatic index was proposed by Ikai (1980) and evaluates the thermostability of proteins based on the percentage of each of the aliphatic amino acids that build up proteins.

Value

The computed aliphatic index for a given amino-acids sequence

References

Ikai (1980). Thermostability and aliphatic index of globular proteins. *Journal of Biochemistry*, 88(6), 1895-1898.

Examples

```
# COMPARED TO EXPASY ALIPHATIC INDEX
# http://web.expasy.org/protparam/
# SEQUENCE: SDKEVDEVDAALSDLEITLE
# Aliphatic index: 117.00

aIndex(seq = "SDKEVDEVDAALSDLEITLE")
# [1] 117
```

autoCorrelation	<i>Compute the auto-correlation index of a protein sequence</i>
-----------------	-----------------------------------------------------------------

Description

This function computes the Cruciani et al (2004) auto-correlation index. The autoCorrelation index is calculated for a lag 'd' using a descriptor 'f' (centred) over a sequence of length 'L'.

Usage

```
autoCorrelation(sequence, lag, property, center = TRUE)
```

Arguments

sequence	An amino-acids sequence
lag	A value for a lag, the max value is equal to the length of shortest peptide minus one.
property	A property to use as value to be correlated.
center	A logical value TRUE or FALSE if the property must be centered.

Value

The computed auto-correlation index for a given amino-acids sequence

References

Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. J. Chemom. 18, 146-155.

Examples

```
# Loading a property to evaluate its autocorrelation
data(AAdata)

# Calculate the auto-correlation index for a lag=1
autoCorrelation(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 1,
  property = AAdata$Hydrophobicity$KyteDoolittle,
  center = TRUE
)
# [1] -0.3519908

# Calculate the auto-correlation index for a lag=5
autoCorrelation(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 5,
```

```

    property = Adata$Hydrophobicity$KyteDoolittle,
    center = TRUE
  )
# [1] 0.001133553

```

autoCovariance *Compute the auto-covariance index of a protein sequence*

Description

This function computes the Cruciani et al (2004) auto-covariance index. The autoCovariance index is calculated for a lag 'd' using a descriptor 'f' (centred) over a sequence of length 'L'.

Usage

```
autoCovariance(sequence, lag, property, center = TRUE)
```

Arguments

sequence	An amino-acids sequence
lag	A value for a lag, the max value is equal to the length of the shortest peptide minus one.
property	A property to use as value to evaluate the covariance.
center	A logical value TRUE or FALSE if the property must be centered.

Value

The computed auto-covariance index for a given amino-acids sequence

References

Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. *J. Chemom.* 18, 146-155.

Examples

```

# Loading a property to evaluate its autocorrelation
data(Adata)

# Calculate the auto-covariance index for a lag=1
autoCovariance(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 1,
  property = Adata$Hydrophobicity$KyteDoolittle,
  center = TRUE
)
# [1] -0.4140053

```

```
# Calculate the auto-covariance index for a lag=5
autoCovariance(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 5,
  property = Aadata$Hydrophobicity$KyteDoolittle,
  center = TRUE
)
# [1] 0.001000336
```

blosumIndices

Compute the BLOSUM62 derived indices of a protein sequence

Description

BLOSUM indices were derived of physicochemical properties that have been subjected to a VARI-MAX analyses and an alignment matrix of the 20 natural AAs using the BLOSUM62 matrix.

Usage

```
blosumIndices(seq)
```

Arguments

seq An amino-acids sequence

Value

The computed average of BLOSUM indices of all the amino acids in the corresponding peptide sequence.

References

Georgiev, A. G. (2009). Interpretable numerical descriptors of amino acid space. *Journal of Computational Biology*, 16(5), 703-723.

Examples

```
blosumIndices(seq = "KLKLLLLLKLK")
# [[1]]
# BLOSUM1 BLOSUM2 BLOSUM3 BLOSUM4 BLOSUM5
# -0.4827273 -0.5618182 -0.8509091 -0.4172727 0.3172727

# BLOSUM6 BLOSUM7 BLOSUM8 BLOSUM9 BLOSUM10
# 0.2527273 0.1463636 0.1427273 -0.2145455 -0.3218182
```

`boman`*Compute the Boman (Potential Protein Interaction) index*

Description

This function computes the potential protein interaction index proposed by Boman (2003) based in the amino acid sequence of a protein. The index is equal to the sum of the solubility values for all residues in a sequence, it might give an overall estimate of the potential of a peptide to bind to membranes or other proteins as receptors, to normalize it is divided by the number of residues. A protein have high binding potential if the index value is higher than 2.48.

Usage`boman(seq)`**Arguments**

<code>seq</code>	An amino-acid sequence
------------------	------------------------

Details

The potential protein interaction index was proposed by Boman (2003) as an easy way to differentiate the action mechanism of hormones (protein-protein) and antimicrobial peptides (protein-membrane) through this index. This function predicts the potential peptide interaction with another protein.

Value

The computed potential protein-protein interaction for a given amino-acids sequence

References

Boman, H. G. (2003). Antibacterial peptides: basic facts and emerging concepts. *Journal of Internal Medicine*, 254(3), 197-215.

Examples

```
# COMPARED TO YADAMP DATABASE
# http://yadamp.unisa.it/showItem.aspx?yadampid=845&x=0,4373912
# SEQUENCE: FLPVLAGLTPSIVPKLVCLLTKKC
# BOMAN INDEX -1.24

boman(seq= "FLPVLAGLTPSIVPKLVCLLTKKC")
# [1] -1.235833
```

charge *Compute the theoretical net charge of a protein sequence*

Description

This function computes the net charge of a protein sequence based on the Henderson-Hasselbalch equation described by Moore, D. S. (1985). The net charge can be calculated at defined pH using one of the 9 pKa scales available: Bjellqvist, Dawson, EMBOSS, Lehninger, Murray, Rodwell, Sillero, Solomon or Stryer.

Usage

```
charge(seq, pH = 7, pKscale = "Lehninger")
```

Arguments

seq	An amino-acids sequence
pH	A pH value
pKscale	A character string specifying the pKa scale to be used; must be one of "Bjellqvist", "Dawson", "EMBOSS", "Lehninger", "Murray", "Rodwell", "Sillero", "Solomon" or "Stryer"

Author(s)

Original by Daniel Osorio <dcosorih@tamu.edu>, C++ code optimized by Luis Pedro Coelho <luis@luispedro.org>

References

- Kiraga, J. (2008) Analysis and computer simulations of variability of isoelectric point of proteins in the proteomes. PhD thesis, University of Wroclaw, Poland.
- Bjellqvist, B., Hughes, G.J., Pasquali, Ch., Paquet, N., Ravier, F., Sanchez, J.Ch., Frutiger S., Hochstrasser D. (1993) The focusing positions of polypeptides in immobilized pH gradients can be predicted from their amino acid sequences. *Electrophoresis*, 14:1023-1031.
- Dawson, R. M. C.; Elliot, D. C.; Elliot, W. H.; Jones, K. M. *Data for biochemical research*. Oxford University Press, 1989; p. 592.
- EMBOSS data are from <http://emboss.sourceforge.net/apps/release/5.0/emboss/apps/iep.html>.
- Nelson, D. L.; Cox, M. M. *Lehninger Principles of Biochemistry*, Fourth Edition; W. H. Freeman, 2004; p. 1100.
- Murray, R.K., Granner, D.K., Rodwell, V.W. (2006) *Harper's illustrated Biochemistry*. 27th edition. Published by The McGraw-Hill Companies.
- Rodwell, J. Heterogeneity of component bands in isoelectric focusing patterns. *Analytical Biochemistry*, 1982, 119 (2), 440-449.
- Sillero, A., Maldonado, A. (2006) Isoelectric point determination of proteins and other macromolecules: oscillating method. *Comput Biol Med.*, 36:157-166.

Solomon, T.W.G. (1998) Fundamentals of Organic Chemistry, 5th edition. Published by Wiley.

Stryer L. (1999) Biochemia. czwarta edycja. Wydawnictwo Naukowe PWN.

Examples

```
# COMPARED TO EMBOSS PEPSTATS
# http://emboss.bioinformatics.nl/cgi-bin/emboss/pepstats
# SEQUENCE: FLPVLAGLTPSIVPKLVCLLTKKC
# Charge = 3.0

charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Bjellqvist")
# [1] 2.737303
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "EMBOSS")
# [1] 2.914112
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Murray")
# [1] 2.907541
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Sillero")
# [1] 2.919812
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Solomon")
# [1] 2.844406
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Stryer")
# [1] 2.876504
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Lehninger")
# [1] 2.87315
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Dawson")
# [1] 2.844406
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "Rodwell")
# [1] 2.819755

# COMPARED TO YADAMP
# http://yadamp.unisa.it/showItem.aspx?yadampid=845&x=0,7055475
# SEQUENCE: FLPVLAGLTPSIVPKLVCLLTKKC
# CHARGE pH5: 3.00
# CHARGE pH7: 2.91
# CHARGE pH9: 1.09
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 5, pKscale= "EMBOSS")
# [1] 3.037398
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 7, pKscale= "EMBOSS")
# [1] 2.914112
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= 9, pKscale= "EMBOSS")
# [1] 0.7184524

# JUST ONE COMMAND
charge(seq= "FLPVLAGLTPSIVPKLVCLLTKKC",pH= seq(from = 5,to = 9,by = 2), pKscale= "EMBOSS")
# [1] 3.0373984 2.9141123 0.7184524
```

Description

This function computes the Cruciani et al (2004) cross-covariance index. The lagged crossCovariance index is calculated for a lag 'd' using two descriptors 'f1' and 'f2' (centred) over a sequence of length 'L'.

Usage

```
crossCovariance(sequence, lag, property1, property2, center = TRUE)
```

Arguments

sequence	An amino-acids sequence
lag	A value for a lag, the max value is equal to the length of the shortest peptide minus one.
property1	A property to use as value to evaluate the cross-covariance.
property2	A property to use as value to evaluate the cross-covariance.
center	A logical value TRUE or FALSE if the property must be centered.

Value

The computed cross-covariance index for a given amino-acids sequence

References

Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. *J. Chemom.* 18, 146-155.

Examples

```
# Loading a property to evaluate its autocorrelation
data(AAdata)

# Calculate the cross-covariance index for a lag=1
crossCovariance(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 1,
  property1 = AAdata$Hydrophobicity$KyteDoolittle,
  property2 = AAdata$Hydrophobicity$Eisenberg,
  center = TRUE
)
# [1] -0.3026609

# Calculate the cross-correlation index for a lag=5
crossCovariance(
  sequence = "SDKEVDEVDAALSDLEITLE",
  lag = 5,
  property1 = AAdata$Hydrophobicity$KyteDoolittle,
  property2 = AAdata$Hydrophobicity$Eisenberg,
```

```
    center = TRUE
  )
# [1] 0.02598035
```

crucianiProperties *Compute the Cruciani properties of a protein sequence*

Description

This function calculates the Cruciani properties of an amino-acids sequence using the scaled principal component scores that summarize a broad set of descriptors calculated based on the interaction of each amino acid residue with several chemical groups (or "probes"), such as charged ions, methyl, hydroxyl groups, and so forth.

Usage

```
crucianiProperties(seq)
```

Arguments

seq An amino-acids sequence

Value

The computed average of Cruciani properties of all the amino acids in the corresponding peptide sequence. Each PP represent an amino-acid property as follows:

- PP1: Polarity,
- PP2: Hydrophobicity,
- PP3: H-bonding

References

Cruciani, G., Baroni, M., Carosati, E., Clementi, M., Valigi, R., and Clementi, S. (2004) Peptide studies by means of principal properties of amino acids derived from MIF descriptors. *J. Chemom.* 18, 146-155.

Examples

```
crucianiProperties(seq = "QWGRCCGWGPGRRYCVRWC")
#   PP1   PP2   PP3
# -0.1130 -0.0220  0.2735
```

`fasgaiVectors`*Compute the FASGAI vectors of a protein sequence*

Description

The FASGAI vectors (Factor Analysis Scales of Generalized Amino Acid Information) is a set of amino acid descriptors, that reflects hydrophobicity, alpha and turn propensities, bulky properties, compositional characteristics, local flexibility, and electronic properties, that can be utilized to represent the sequence structural features of peptides or protein motifs.

Usage

```
fasgaiVectors(seq)
```

Arguments

<code>seq</code>	An amino-acids sequence
------------------	-------------------------

Value

The computed average of FASGAI factors of all the amino acids in the corresponding peptide sequence. Each factor represent an amino-acid property as follows:

- F1: Hydrophobicity index,
- F2: Alpha and turn propensities,
- F3: Bulky properties,
- F4: Compositional characteristic index,
- F5: Local flexibility,
- F6: Electronic properties

References

Liang, G., & Li, Z. (2007). Factor analysis scale of generalized amino acid information as the source of a new set of descriptors for elucidating the structure and activity relationships of cationic antimicrobial peptides. *Molecular Informatics*, 26(6), 754-763.

Examples

```
fasgaiVectors(seq = "QWGRRC CGWGPGRRYCVRWC")
# [[1]]
#   F1      F2      F3      F4      F5      F6
# -0.13675 -0.45485 -0.11695 -0.45800 -0.38015  0.52740
```

hmoment

Compute the hydrophobic moment of a protein sequence

Description

This function compute the hmoment based on Eisenberg, D., Weiss, R. M., & Terwilliger, T. C. (1984). Hydrophobic moment is a quantitative measure of the amphiphilicity perpendicular to the axis of any periodic peptide structure, such as the a-helix or b-sheet. It can be calculated for an amino acid sequence of N residues and their associated hydrophobicities H_n .

Usage

```
hmoment(seq, angle = 100, window = 11)
```

Arguments

seq	An amino-acids sequence
angle	A protein rotational angle (Suggested: a-helix = 100, b-sheet=160)
window	A sequence fraction length

Details

The hydrophobic moment was proposed by Eisenberg et al. (1982), as a quantitative measure of the amphiphilicity perpendicular to the axis of any periodic peptide structure. It is computed using the standardized Eisenberg (1984) scale, windows (fragment of sequence) of eleven amino acids (by default) and specifying the rotational angle at which it should be calculated.

Value

The computed maximal hydrophobic moment (uH) for a given amino-acids sequence

Note

This function was written by an anonymous reviewer of the RJournal

References

Eisenberg, D., Weiss, R. M., & Terwilliger, T. C. (1984). The hydrophobic moment detects periodicity in protein hydrophobicity. *Proceedings of the National Academy of Sciences*, 81(1), 140-144.

Examples

```
# COMPARED TO EMBOSS:HMOMENT
# http://emboss.bioinformatics.nl/cgi-bin/emboss/hmoment
# SEQUENCE: FLPVLAGLTPSIVPKLVCLLTKKC
# ALPHA-HELIX ANGLE=100 : 0.52
# BETA-SHEET ANGLE=160 : 0.271
```

```
# ALPHA HELIX VALUE
hmoment(seq = "FLPVLAGLTPSIVPKLVCLLTKKC", angle = 100, window = 11)
# [1] 0.5199226

# BETA SHEET VALUE
hmoment(seq = "FLPVLAGLTPSIVPKLVCLLTKKC", angle = 160, window = 11)
# [1] 0.2705906
```

hydrophobicity

Compute the hydrophobicity index of a protein sequence

Description

This function calculates the GRAVY hydrophobicity index of an amino acids sequence using one of the 38 scales from different sources.

Usage

```
hydrophobicity(seq, scale = "KyteDoolittle")
```

Arguments

seq	An amino-acids sequence
scale	A character string specifying the hydrophobicity scale to be used; must be one of "Aboderin", "AbrahamLeo", "Argos", "BlackMould", "BullBreese", "Casari", "Chothia", "Cid", "Cowan3.4", "Cowan7.5", "Eisenberg", "Engelman", "Fasman", "Fauchere", "Goldsack", "Guy", "HoppWoods", "Janin", "Jones", "Juretic", "Kidera", "Kuhn", "KyteDoolittle", "Levitt", "Manavalan", "Miyazawa", "Parker", "Ponnuswamy", "Prabhakaran", "Rao", "Rose", "Roseman", "Sweet", "Tanford", "Welling", "Wilson", "Wolfenden", "Zimmerman", "interfaceScale_pH8", "interfaceScale_pH2", "octanolScale_pH8", "octanolScale_pH2", "oiScale_pH8" or "oiScale_pH2".

Details

The hydrophobicity is an important stabilization force in protein folding; this force changes depending on the solvent in which the protein is found. The hydrophobicity index is calculated adding the hydrophobicity of individual amino acids and dividing this value by the length of the sequence.

Value

The computed GRAVY index for a given amino-acid sequence

References

- Aboderin, A. A. (1971). An empirical hydrophobicity scale for alpha-amino-acids and some of its applications. *International Journal of Biochemistry*, 2(11), 537-544.
- Abraham D.J., Leo A.J. Hydrophobicity ($\Delta G_{1/2}$ cal). *Proteins: Structure, Function and Genetics* 2:130-152(1987).
- Argos, P., Rao, J. K., & Hargrave, P. A. (1982). Structural Prediction of Membrane-Bound Proteins. *European Journal of Biochemistry*, 128(2-3), 565-575.
- Black S.D., Mould D.R. Hydrophobicity of physiological L-alpha amino acids. *Anal. Biochem.* 193:72-82(1991).
- Bull H.B., Breese K. Hydrophobicity (free energy of transfer to surface in kcal/mole). *Arch. Biochem. Biophys.* 161:665-670(1974).
- Casari, G., & Sippl, M. J. (1992). Structure-derived hydrophobic potential: hydrophobic potential derived from X-ray structures of globular proteins is able to identify native folds. *Journal of molecular biology*, 224(3), 725-732.
- Chothia, C. (1976). The nature of the accessible and buried surfaces in proteins. *Journal of molecular biology*, 105(1), 1-12.
- Cid, H., Bunster, M., Canales, M., & Gazitua, F. (1992). Hydrophobicity and structural classes in proteins. *Protein engineering*, 5(5), 373-375.
- Cowan R., Whittaker R.G. Hydrophobicity indices at pH 3.4 determined by HPLC. *Peptide Research* 3:75-80(1990).
- Cowan R., Whittaker R.G. Hydrophobicity indices at pH 7.5 determined by HPLC. *Peptide Research* 3:75-80(1990).
- Eisenberg D., Schwarz E., Komarony M., Wall R. Normalized consensus hydrophobicity scale. *J. Mol. Biol.* 179:125-142(1984).
- Engelman, D. M., Steitz, T. A., & Goldman, A. (1986). Identifying nonpolar transbilayer helices in amino acid sequences of membrane proteins. *Annual review of biophysics and biophysical chemistry*, 15(1), 321-353.
- Fasman, G. D. (Ed.). (1989). *Prediction of protein structure and the principles of protein conformation*. Springer.
- Fauchere J.-L., Pliska V.E. Hydrophobicity scale (π - r). *Eur. J. Med. Chem.* 18:369-375(1983).
- Goldsack, D. E., & Chalifoux, R. C. (1973). Contribution of the free energy of mixing of hydrophobic side chains to the stability of the tertiary structure of proteins. *Journal of theoretical biology*, 39(3), 645-651.
- Guy H.R. Hydrophobicity scale based on free energy of transfer (kcal/mole). *Biophys J.* 47:61-70(1985).
- Hopp T.P., Woods K.R. Hydrophilicity. *Proc. Natl. Acad. Sci. U.S.A.* 78:3824-3828(1981).
- Janin J. Free energy of transfer from inside to outside of a globular protein. *Nature* 277:491-492(1979).
- Jones, D. D. (1975). Amino acid properties and side-chain orientation in proteins: a cross correlation approach. *Journal of theoretical biology*, 50(1), 167-183.
- Juretic, D., Lucic, B., Zucic, D., & Trinajstic, N. (1998). Protein transmembrane structure: recognition and prediction by using hydrophobicity scales through preference functions. *Theoretical and computational chemistry*, 5, 405-445.

- Kidera, A., Konishi, Y., Oka, M., Ooi, T., & Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, 4(1), 23-55.
- Kuhn, L. A., Swanson, C. A., Pique, M. E., Tainer, J. A., & Getzoff, E. D. (1995). Atomic and residue hydrophilicity in the context of folded protein structures. *Proteins: Structure, Function, and Bioinformatics*, 23(4), 536-547.
- Kyte J., Doolittle R.F. Hydrophobicity. *J. Mol. Biol.* 157:105-132(1982).
- Levitt, M. (1976). A simplified representation of protein conformations for rapid simulation of protein folding. *Journal of molecular biology*, 104(1), 59-107.
- Manavalan P., Ponnuswamy Average surrounding hydrophobicity. P.K. *Nature* 275:673-674(1978).
- Miyazawa S., Jernigen R.L. Hydrophobicity scale (contact energy derived from 3D data). *Macromolecules* 18:534-552(1985).
- Parker J.M.R., Guo D., Hodges R.S. Hydrophilicity scale derived from HPLC peptide retention times. *Biochemistry* 25:5425-5431(1986).
- Ponnuswamy, P. K. (1993). Hydrophobic characteristics of folded proteins. *Progress in biophysics and molecular biology*, 59(1), 57-103.
- Prabhakaran, M. (1990). The distribution of physical, chemical and conformational properties in signal and nascent peptides. *Biochem. J.*, 269, 691-696.
- Rao M.J.K., Argos P. Membrane buried helix parameter. *Biochim. Biophys. Acta* 869:197-214(1986).
- Rose G.D., Geselowitz A.R., Lesser G.J., Lee R.H., Zehfus M.H. Mean fractional area loss (f) [average area buried/standard state area]. *Science* 229:834-838(1985)
- Roseman M.A. Hydrophobicity scale (pi-r). *J. Mol. Biol.* 200:513-522(1988).
- Sweet R.M., Eisenberg D. Optimized matching hydrophobicity (OMH). *J. Mol. Biol.* 171:479-488(1983).
- Tanford C. Hydrophobicity scale (Contribution of hydrophobic interactions to the stability of the globular conformation of proteins). *J. Am. Chem. Soc.* 84:4240-4274(1962).
- Welling G.W., Weijer W.J., Van der Zee R., Welling-Wester S. Antigenicity value X 10. *FEBS Lett.* 188:215-218(1985).
- Wilson K.J., Honegger A., Stotzel R.P., Hughes G.J. Hydrophobic constants derived from HPLC peptide retention times. *Biochem. J.* 199:31-41(1981).
- Wolfenden R.V., Andersson L., Cullis P.M., Southgate C.C.F. Hydration potential (kcal/mole) at 25C. *Biochemistry* 20:849-855(1981).
- Zimmerman, J. M., Eliezer, N., & Simha, R. (1968). The characterization of amino acid sequences in proteins by statistical methods. *Journal of theoretical biology*, 21(2), 170-201.
- Nakai, K., Kidera, A., and Kanehisa, M.; Cluster analysis of amino acid indices for prediction of protein structure and function. *Protein Eng.* 2, 93-100 (1988).
- Tomii, K. and Kanehisa, M.; Analysis of amino acid indices and mutation matrices for sequence comparison and structure prediction of proteins. *Protein Eng.* 9, 27-36 (1996).
- Kawashima, S., Ogata, H., and Kanehisa, M.; AAindex: amino acid index database. *Nucleic Acids Res.* 27, 368-369 (1999).

Kawashima, S. and Kanehisa, M.; AAindex: amino acid index database. *Nucleic Acids Res.* 28, 374 (2000).

Kawashima, S., Pokarowski, P., Pokarowska, M., Kolinski, A., Katayama, T., and Kanehisa, M.; AAindex: amino acid index database, progress report 2008. *Nucleic Acids Res.* 36, D202-D205 (2008).

White, Stephen (2006-06-29). "Experimentally Determined Hydrophobicity Scales". University of California, Irvine. Retrieved 2017-05-25

Examples

```
# COMPARED TO GRAVY Grand average of hydropathicity (GRAVY) ExpASY
# http://web.expasy.org/cgi-bin/protparam/protparam
# SEQUENCE: QWGRCCGWGPGRRYCVRWC
# GRAVY: -0.950

hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Aboderin")
#[1] 3.84
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "AbrahamLeo")
#[1] 0.092
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Argos")
#[1] 1.033
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "BlackMould")
#[1] 0.50125
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "BullBreese")
#[1] 0.1575
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Casari")
#[1] 0.38
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Chothia")
#[1] 0.262
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Cid")
#[1] 0.198
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Cowan3.4")
#[1] 0.0845
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Cowan7.5")
#[1] 0.0605
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Eisenberg")
#[1] -0.3265
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Engelman")
#[1] 2.31
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Fasman")
#[1] -1.2905
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Fauchere")
#[1] 0.527
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Goldsack")
#[1] 1.2245
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Guy")
#[1] 0.193
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "HoppWoods")
#[1] -0.14
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Janin")
#[1] -0.105
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC", scale = "Jones")
```

```

#[1] 1.4675
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Juretic")
#[1] -1.106
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Kidera")
#[1] -0.0405
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Kuhn")
#[1] 0.9155
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "KyteDoolittle")
#[1] -0.95
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Levitt")
#[1] -0.21
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Manavalan")
#[1] 13.0445
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Miyazawa")
#[1] 5.739
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Parker")
#[1] 1.095
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Ponnuswamy")
#[1] 0.851
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Prabhakaran")
#[1] 9.67
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Rao")
#[1] 0.813
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Rose")
#[1] 0.7575
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Roseman")
#[1] -0.495
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Sweet")
#[1] -0.1135
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Tanford")
#[1] -0.2905
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Welling")
#[1] -0.666
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Wilson")
#[1] 3.16
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Wolfenden")
#[1] -6.307
hydrophobicity(seq = "QWGRCCGWGPGRRYCVRWC",scale = "Zimmerman")
#[1] 0.943

```

Description

This function calculates the instability index proposed by Guruprasad (1990). This index predicts the stability of a protein based on its amino acid composition, a protein whose instability index is smaller than 40 is predicted as stable, a value above 40 predicts that the protein may be unstable.

Usage

```
instaIndex(seq)
```

Arguments

```
seq          An amino-acids sequence
```

Value

The computed instability index for a given amino-acids sequence

References

Guruprasad K, Reddy BV, Pandit MW (1990). "Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence". *Protein Eng.* 4 (2): 155 - 61. doi:10.1093/protein/4.2.155

Examples

```
# COMPARED TO ExPASy INSTAINDEX
# http://web.expasy.org/protparam/
# SEQUENCE: QWGRCCGWGPGRRYCVRWC
# The instability index (II) is computed to be 83.68

instaIndex(seq = "QWGRCCGWGPGRRYCVRWC")
# [1] 83.68
```

kideraFactors

Compute the Kidera factors of a protein sequence

Description

The Kidera Factors were originally derived by applying multivariate analysis to 188 physical properties of the 20 amino acids and using dimension reduction techniques. This function calculates the average of the ten Kidera factors for a protein sequence.

Usage

```
kideraFactors(seq)
```

Arguments

```
seq          An amino-acids sequence
```

Value

A list with the average of the ten Kidera factors. The first four factors are essentially pure physical properties; the remaining six factors are superpositions of several physical properties, and are labelled for convenience by the name of the most heavily weighted component.

- KF1: Helix/bend preference,
- KF2: Side-chain size,
- KF3: Extended structure preference,
- KF4: Hydrophobicity,
- KF5: Double-bend preference,
- KF6: Partial specific volume,
- KF7: Flat extended preference,
- KF8: Occurrence in alpha region,
- KF9: pK-C,
- KF10: Surrounding hydrophobicity

References

Kidera, A., Konishi, Y., Oka, M., Ooi, T., & Scheraga, H. A. (1985). Statistical analysis of the physical properties of the 20 naturally occurring amino acids. *Journal of Protein Chemistry*, 4(1), 23-55.

Examples

```
kideraFactors(seq = "KLKLLLLLKLK")
# [[1]]
#   KF1      KF2      KF3      KF4      KF5
# -0.78545455 0.29818182 -0.23636364 -0.08181818 0.21000000
#   KF6      KF7      KF8      KF9      KF10
# -1.89363636 1.02909091 -0.51272727 0.11181818 0.81000000
```

lengthpep

Compute the amino acid length of a protein sequence

Description

This function counts the number of amino acids in a protein sequence

Usage

```
lengthpep(seq)
```

Arguments

seq An amino-acids sequence

Details

All proteins are formed by linear chains of small residues known as amino acids attached to each other by peptide bonds. The function `lengthpep` counts the number of amino acids in a sequence and returns a vector with the count for each peptide used as argument.

Examples

```
# COMPARED TO ExPASy ProtParam
# http://web.expasy.org/protparam
# SEQUENCE: QWGRCCGWGPGRRYCVRWC
# Number of amino acids: 20

lengthpep(seq = "QWGRCCGWGPGRRYCVRWC")
# [1] 20
```

massShift

Calculate the mass difference of modified peptides.

Description

This function calculates the mass difference of peptides introduced by chemical modifications or heavy isotope labelling.

Usage

```
massShift(seq, label = "none", aaShift = NULL, monoisotopic = TRUE)
```

Arguments

<code>seq</code>	An amino-acids sequence, in one letter code.
<code>label</code>	Set a predefined heavy isotope label. Accepts "none", "silac_13c", "silac_13c15n" and "15n". Overwrites input in <code>aaShift</code> .
<code>aaShift</code>	Define the mass difference in Dalton of given amino acids as a named vector. Use the amino acid one letter code as names and the mass shift in Dalton as values.
<code>monoisotopic</code>	A logical value 'TRUE' or 'FALSE' indicating if monoisotopic weights of amino-acids should be used

Source

For the predefined heavy isotope labels, compare:

- silac_13c [Unimod 188](#)
- silac_13c15n [Unimod 259](#) and [Unimod 267](#)
- 15n [Unimod 994](#), [Unimod 995](#), [Unimod 996](#) and [Unimod 897](#)

Examples

```
massShift("EGVNDNECEGFFSAR", label = "silac_13c")
massShift("EGVNDNECEGFFSAR", aaShift = c(K = 6.020129, R = 6.020129))
```

membpos

Compute theoretically the class of a protein sequence

Description

This function calculates the theoretical class of a protein sequence based on the relationship between the hydrophobic moment and hydrophobicity scale proposed by Eisenberg (1984).

Usage

```
membpos(seq, angle = 100)
```

Arguments

seq	An amino-acids sequence
angle	A protein rotational angle

Details

Eisenberg et al. (1982) found a correlation between hydrophobicity and hydrophobic moment that defines the protein section as globular, transmembrane or superficial. The function calculates the hydrophobicity (H) and hydrophobic moment (uH) based on the standardized scale of Eisenberg (1984) using windows of 11 amino acids for calculate the theoretical fragment type.

Value

A data frame for each sequence given with the calculated class for each window of eleven amino-acids

References

Eisenberg, David. "Three-dimensional structure of membrane and surface proteins." Annual review of biochemistry 53.1 (1984): 595-623.

D. Eisenberg, R. M. Weiss, and T. C. Terwilliger. The helical hydrophobic moment: A measure of the amphiphilicity of a helix. Nature, 299(5881):371-374, 1982. [p7, 8]

Examples

```
membpos(seq = "ARQQNLFINFCLILIFLLLI", angle = 100)
#      Pep      H      uH      MembPos
# 1 ARQQNLFINFCL 0.083 0.353      Globular
# 2 RQQNLFINFCLI 0.147 0.317      Globular
# 3 QQNLFINFCLIL 0.446 0.274      Globular
# 4 QNLFINFCLILI 0.632 0.274 Transmembrane
# 5 NLFINFCLILIF 0.802 0.253      Surface
# 6 LFINFCLILIFL 0.955 0.113 Transmembrane
# 7 FINFCLILIFLL 0.955 0.113 Transmembrane
# 8 INFCLILIFLLL 0.944 0.108 Transmembrane
# 9 NFCLILIFLLLI 0.944 0.132 Transmembrane
```

```
membpos(seq = "ARQQNLFINFCLILIFLLLI", angle = 160)
#      Pep      H      uH      MembPos
# 1 ARQQNLFINFCL 0.083 0.467      Globular
# 2 RQQNLFINFCLI 0.147 0.467      Globular
# 3 QQNLFINFCLIL 0.446 0.285      Globular
# 4 QNLFINFCLILI 0.632 0.358      Surface
# 5 NLFINFCLILIF 0.802 0.358      Surface
# 6 LFINFCLILIFL 0.955 0.269      Surface
# 7 FINFCLILIFLL 0.955 0.269      Surface
# 8 INFCLILIFLLL 0.944 0.257      Surface
# 9 NFCLILIFLLLI 0.944 0.229      Surface
```

mswhimScores

Compute the MS-WHIM scores of a protein sequence

Description

MS-WHIM scores were derived from 36 electrostatic potential properties derived from the three-dimensional structure of the 20 natural amino acids

Usage

```
mswhimScores(seq)
```

Arguments

```
seq          An amino-acids sequence
```

Value

The computed average of MS-WHIM scores of all the amino acids in the corresponding peptide sequence.

References

Zaliani, A., & Gancia, E. (1999). MS-WHIM scores for amino acids: a new 3D-description for peptide QSAR and QSPR studies. *Journal of chemical information and computer sciences*, 39(3), 525-533.

Examples

```
mswhimScores(seq = "KLKLLLLLKLK")
# [[1]]
# MSWHIM1 MSWHIM2 MSWHIM3
# -0.6563636 0.4872727 0.1163636
```

mw

Compute the molecular weight of a protein sequence

Description

This function calculates the molecular weight of a protein sequence. It is calculated as the sum of the mass of each amino acid using the scale available on Compute pI/Mw tool. It also supports mass calculation of proteins with predefined or custom stable isotope mass labels.

Usage

```
mw(seq, monoisotopic = FALSE, label = "none", aaShift = NULL)
```

Arguments

seq	An amino-acids sequence
monoisotopic	A logical value 'TRUE' or 'FALSE' indicating if monoisotopic weights of amino-acids should be used
label	Set a predefined heavy isotope label. Accepts "none", "silac_13c", "silac_13c15n" and "15n". Overwrites input in aaShift.
aaShift	Define the mass difference in Dalton of given amino acids as a named vector. Use the amino acid one letter code as names and the mass shift in Dalton as values.

Details

The molecular weight is the sum of the masses of each atom constituting a molecule. The molecular weight is directly related to the length of the amino acid sequence and is expressed in units called daltons (Da). In Peptides the function mw computes the molecular weight using the same formulas and weights as ExPASy's "compute pI/mw" tool (Gasteiger et al., 2005).

Source

The formula and amino acid scale are the same available on ExPASy Compute pI/Mw tool: http://web.expasy.org/compute_pi

References

Gasteiger, E., Hoogland, C., Gattiker, A., Wilkins, M. R., Appel, R. D., & Bairoch, A. (2005). Protein identification and analysis tools on the ExPASy server. In *The proteomics protocols handbook* (pp. 571-607). Humana Press. Chicago

Examples

```
# COMPARED TO ExPASy Compute pI/Mw tool
# http://web.expasy.org/compute_pi/
# SEQUENCE: QWGRRC CGWGPGRRYCVRWC
# Theoretical pI/Mw: 9.88 / 2485.91

mw(seq = "QWGRRC CGWGPGRRYCVRWC", monoisotopic = FALSE)
# [1] 2485.911

mw(seq = "QWGRRC CGWGPGRRYCVRWC", monoisotopic = TRUE)
# [1] 2484.12
```

mz

Calculate the m/z for peptides.

Description

This function calculates the (monoisotopic) mass over charge ratio (m/z) for peptides, as measured in mass spectrometry.

Usage

```
mz(seq, charge = 2, label = "none", aaShift = NULL, cysteins = 57.021464)
```

Arguments

seq	An amino-acids sequence, in one letter code.
charge	The net charge for which the m/z should be calculated
label	Set a predefined heavy isotope label. Accepts "none", "silac_13c", "silac_13c15n" and "15n". Overwrites input in aaShift.
aaShift	Define the mass difference in Dalton of given amino acids as a named vector. Use the amino acid one letter code as names and the mass shift in Dalton as values.
cysteins	Define the mass shift in Dalton of blocked cysteins. Defaults to 57.021464, for cysteins blocked by iodoacetamide.

Examples

```
mz("EGVNDNECEGFFSAR")
mz("EGVNDNECEGFFSAR", aaShift = c(K = 6.020129, R = 6.020129))
mz("EGVNDNECEGFFSAR", label = "silac_13c", cysteins = 58.005479)
```

pepdata	<i>Physicochemical properties and indices from 100 amino acid sequences</i>
---------	-----------------------------------------------------------------------------

Description

Physicochemical properties and indices from 100 amino acid sequences (50 antimicrobial and 50 non antimicrobial)

Usage

```
data(pepdata)
```

Format

A data frame with 100 observations on the following 23 variables.

sequence a character vector with the sequences of 100 peptides (50 antimicrobial and 50 non-antimicrobial)

group Integer vector with the group code "0" for non antimicrobial and "1" for antimicrobial

length a numeric vector with the length of the amino acid sequence

mw a numeric vector with the molecular weight of the amino acid sequence

tinyAA A numeric vector with the fraction (as percent) of tiny amino acids that make up the sequence

smallAA A numeric vector with the fraction (as percent) of small amino acids that make up the sequence

aliphaticAA A numeric vector with the fraction (as percent) of aliphatic amino acids that make up the sequence

aromaticAA A numeric vector with the fraction (as percent) of aromatic amino acids that make up the sequence

nonpolarAA A numeric vector with the fraction (as percent) of non-polar amino acids that make up the sequence

polarAA A numeric vector with the fraction (as percent) of polar amino acids that make up the sequence

chargedAA A numeric vector with the fraction (as percent) of charged amino acids that make up the sequence

basicAA A numeric vector with the fraction (as percent) of basic amino acids that make up the sequence

acidicAA A numeric vector with the fraction (as percent) of acid amino acids that make up the sequence

charge a numeric vector with the charge of the amino acid sequence

pI a numeric vector with the isoelectric point of the amino acid sequence

aindex a numeric vector with the aliphatic index of the amino acid sequence

instaindex a numeric vector with the instability index of the amino acid sequence
 boman a numeric vector with the potential peptide-interaction index of the amino acid sequence
 hydrophobicity a numeric vector with the hydrophobicity index of the amino acid sequence
 hmoment a numeric vector with the hydrophobic moment of the amino acid sequence
 transmembrane A numeric vector with the fraction of Transmembrane windows of 11 amino acids that make up the sequence
 surface A numeric vector with the fraction of Surface windows of 11 amino acids that make up the sequence
 globular A numeric vector with the fraction of Globular windows of 11 amino acids that make up the sequence

pI *Compute the isoelectric point (pI) of a protein sequence*

Description

The isoelectric point (pI), is the pH at which a particular molecule or surface carries no net electrical charge.

Usage

```
pI(seq, pKscale = "EMBOSS")
```

Arguments

seq	An amino-acids sequence
pKscale	A character string specifying the pK scale to be used; must be one of "Bjellqvist", "EMBOSS", "Murray", "Sillero", "Solomon", "Stryer", "Lehninger", "Dawson" or "Rodwell"

Details

The isoelectric point (pI) is the pH at which the net charge of the protein is equal to 0. It is a variable that affects the solubility of the peptides under certain conditions of pH. When the pH of the solvent is equal to the pI of the protein, it tends to precipitate and lose its biological function.

Examples

```

# COMPARED TO ExpASy ProtParam
# http://web.expasy.org/cgi-bin/protparam/protparam
# SEQUENCE: QWGRCCGWGPGRRYCVRWC
# Theoretical pI: 9.88

pI(seq= "QWGRCCGWGPGRRYCVRWC",pKscale= "Bjellqvist")
# [1] 9.881

```

```

# COMPARED TO EMBOSS PEPSTATS
# http://emboss.bioinformatics.nl/cgi-bin/emboss/pepstats
# SEQUENCE: QWGRRC CGWGP GRRY CVRWC
# Isoelectric Point = 9.7158

pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "EMBOSS")
# [1] 9.716

# OTHER SCALES

pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Murray")
# [1] 9.818
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Sillero")
# [1] 9.891
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Solomon")
# [1] 9.582
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Stryer")
# [1] 9.623
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Lehninger")
# [1] 9.931
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Dawson")
# [1] 9.568
pI(seq= "QWGRRC CGWGP GRRY CVRWC", pKscale= "Rodwell")
# [1] 9.718

```

plotXVG

Plot time series from GROMACS XVG files

Description

Read and plot output data from a XVG format file.

Usage

```
plotXVG(XVGfile, ...)
```

Arguments

XVGfile	A .XVG output file of the GROMACS molecular dynamics package
...	Arguments to be passed to methods, such as graphical parameters.

Details

GROMACS (GRONingen MACHine for Chemical Simulations) is a molecular dynamics package designed for simulations of proteins, lipids and nucleic acids. It is free, open source software released under the GNU General Public License. The file format used by GROMACS is XVG. This format can be displayed in graphical form through the GRACE program on UNIX/LINUX systems and the GNUPlot program on Windows. XVG files are plain text files containing tabular data separated by tabulators and two types of comments which contain data labels. Although manual

editing is possible, this is not a viable option when working with multiple files of this type. For ease of reading, information management and data plotting, the functions `read.xvg` and `plot.xvg` were incorporated.

Author(s)

Latest: J. Sebastian Paez <jpaezpa@purdue.edu>

Original: Daniel Osorio <dcosorih@unal.edu.co>

References

Pronk, S., Pall, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., ... & Lindahl, E. (2013). GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics*, 29 (7), 845-854.

Examples

```
XVGfile <- system.file("xvg-files/epot.xvg", package="Peptides")
plotXVG(XVGfile)
```

protFP

Compute the protFP descriptors of a protein sequence

Description

The ProtFP descriptor set was constructed from a large initial selection of indices obtained from the AAindex database for all 20 naturally occurring amino acids.

Usage

```
protFP(seq)
```

Arguments

seq An amino-acids sequence

Value

The computed average of protFP descriptors of all the amino acids in the corresponding peptide sequence.

References

van Westen, G. J., Swier, R. F., Wegner, J. K., IJzerman, A. P., van Vlijmen, H. W., & Bender, A. (2013). Benchmarking of protein descriptor sets in proteochemometric modeling (part 1): comparative study of 13 amino acid descriptor sets. *Journal of cheminformatics*, 5(1), 41.

Examples

```
protFP(seq = "QWGRCCGWGPGRRYCVRWC")
# [[1]]
# ProtFP1 ProtFP2 ProtFP3 ProtFP4 ProtFP5 ProtFP6 ProtFP7 ProtFP8
# 0.2065 -0.0565 1.9930 -0.2845 0.7315 0.7000 0.1715 0.1135
```

readXVG	<i>Read output data from a XVG format file.</i>
---------	-------------------------------------------------

Description

XVG is the default format file of the GROMACS molecular dynamics package, contains data formatted to be imported into the Grace 2-D plotting program.

Usage

```
readXVG(file)
```

Arguments

file A .XVG output file of the GROMACS molecular dynamics package

Details

GROMACS (GRONingen MACHine for Chemical Simulations) is a molecular dynamics package designed for simulations of proteins, lipids and nucleic acids. It is free, open source software released under the GNU General Public License. The file format used by GROMACS is XVG. This format can be displayed in graphical form through the GRACE program on UNIX/LINUX systems and the GNUPlot program on Windows. XVG files are plain text files containing tabular data separated by tabulators and two types of comments which contain data labels. Although manual editing is possible, this is not a viable option when working with multiple files of this type. For ease of reading, information management and data plotting, the functions `read.xvg` and `plot.xvg` were incorporated.

Author(s)

Latest: J. Sebastian Paez <jpaezpa@purdue.edu> and hongbo-zhu-cn <@github>

Original: Daniel Osorio <dcosorih@unal.edu.co>

References

Pronk, S., Pall, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., ... & Lindahl, E. (2013). GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics*, 29 (7), 845-854.

Examples

```
# READING FILE
XVGfile <- system.file("xvg-files/epot.xvg",package="Peptides")
readXVG(XVGfile)

#   Time (ps) Potential
# 1      1 6672471040
# 2      2 6516461568
# 3      3 6351947264
# 4      4 6183133184
# 5      5 6015310336
# 6      6 5854271488
```

stScales

Compute the ST-scales of a protein sequence

Description

ST-scales were proposed by Yang et al, taking 827 properties into account which are mainly constitutional, topological, geometrical, hydrophobic, electronic, and steric properties of a total set of 167 AAs.

Usage

```
stScales(seq)
```

Arguments

```
seq           An amino-acids sequence
```

Value

The computed average of ST-scales of all the amino acids in the corresponding peptide sequence.

References

Yang, L., Shu, M., Ma, K., Mei, H., Jiang, Y., & Li, Z. (2010). ST-scale as a novel amino acid descriptor and its application in QSAM of peptides and analogues. *Amino acids*, 38(3), 805-816.

Examples

```
stScales(seq = "QWGRCCGWGPGRRYCVRWC")
# [[1]]
# ST1      ST2      ST3      ST4      ST5      ST6      ST7      ST8
# -0.63760  0.07965  0.05150  0.07135 -0.27905 -0.80995  0.58020  0.54400
```

tScales	<i>Compute the T-scales of a protein sequence</i>
---------	---------------------------------------------------

Description

T-scales are based on 67 common topological descriptors of 135 amino acids. These topological descriptors are based on the connectivity table of amino acids alone, and do not explicitly consider 3D properties of each structure.

Usage

```
tScales(seq)
```

Arguments

seq	An amino-acids sequence
-----	-------------------------

Value

The computed average of T-scales of all the amino acids in the corresponding peptide sequence.

References

Tian F, Zhou P, Li Z: T-scale as a novel vector of topological descriptors for amino acids and its application in QSARs of peptides. *J Mol Struct.* 2007, 830: 106-115. 10.1016/j.molstruc.2006.07.004.

Examples

```
tScales(seq = "QWGRCCGWGPGRRYCVRWC")
# [[1]]
# T1      T2      T3      T4      T5
# -3.2700 -0.0035 -0.3855 -0.1475  0.7585
```

vhseScales	<i>Compute the VHSE-scales of a protein sequence</i>
------------	------------------------------------------------------

Description

VHSE-scales (principal components score Vectors of Hydrophobic, Steric, and Electronic properties), is derived from principal components analysis (PCA) on independent families of 18 hydrophobic properties, 17 steric properties, and 15 electronic properties, respectively, which are included in total 50 physicochemical variables of 20 coded amino acids.

Usage

```
vhseScales(seq)
```

Arguments

seq An amino-acids sequence

Value

The computed average of VHSE-scales of all the amino acids in the corresponding peptide sequence. Each VSHE-scale represent an amino-acid property as follows:

- VHSE1 and VHSE2: Hydrophobic properties
- VHSE3 and VHSE4: Steric properties
- VHSE5 to VHSE8: Electronic properties

References

Mei, H. U., Liao, Z. H., Zhou, Y., & Li, S. Z. (2005). A new set of amino acid descriptors and its application in peptide QSARs. *Peptide Science*, 80(6), 775-786.

Examples

```
vhseScales(seq = "QWGRCCGWGPGRRYCVRWC")
# [[1]]
#  VHSE1  VHSE2  VHSE3  VHSE4  VHSE5  VHSE6  VHSE7  VHSE8
#-0.1150  0.0630 -0.0055  0.7955  0.4355  0.2485  0.1740 -0.0960
```

zScales

Compute the Z-scales of a protein sequence

Description

Z-scales are based on physicochemical properties of the AAs including NMR data and thin-layer chromatography (TLC) data.

Usage

zScales(seq)

Arguments

seq An amino-acids sequence

Value

The computed average of Z-scales of all the amino acids in the corresponding peptide sequence. Each Z scale represent an amino-acid property as follows:

- Z1: Lipophilicity
- Z2: Steric properties (Steric bulk/Polarizability)
- Z3: Electronic properties (Polarity / Charge)
- Z4 and Z5: They relate electronegativity, heat of formation, electrophilicity and hardness.

References

Sandberg M, Eriksson L, Jonsson J, Sjostrom M, Wold S: New chemical descriptors relevant for the design of biologically active peptides. A multivariate characterization of 87 amino acids. J Med Chem 1998, 41:2481-2491.

Examples

```
zScales(seq = "QWGRCCGWGPGRRYCVRWC")
# [[1]]
# Z1      Z2      Z3      Z4      Z5
# 0.6200  0.0865  0.0665  0.7280 -0.8740
```

Index

aaComp, [2](#)
AAdata, [4](#)
aaDescriptors, [9](#)
aaList, [10](#)
aaSMILES, [11](#)
aIndex, [12](#)
autoCorrelation, [13](#)
autoCovariance, [14](#)

blosumIndices, [15](#)
boman, [16](#)

charge, [17](#)
crossCovariance, [18](#)
crucianiProperties, [20](#)

fasgaiVectors, [21](#)

hmoment, [22](#)
hydrophobicity, [23](#)

instaIndex, [27](#)

kideraFactors, [28](#)

lengthpep, [29](#)

massShift, [30](#)
membpos, [31](#)
mswhimScores, [32](#)
mw, [33](#)
mz, [34](#)

pepdata, [35](#)
pI, [36](#)
plotXVG, [37](#)
protFP, [38](#)

readXVG, [39](#)

stScales, [40](#)

tScales, [41](#)
vhseScales, [41](#)

zScales, [42](#)